

Dominions in Filtral Quasivarieties

Miguel Campercholi

Facultad de Matemática, Astronomía y Física
Universidad Nacional de Córdoba
Argentina

Logic Colloquium 2019 - Prague

Dominions

Let $\mathbf{A} \leq \mathbf{B}$ and \mathcal{K} a class. Say that $b \in \mathbf{B}$ is **dominated** by \mathbf{A} w.r.t. \mathcal{K} if

Dominions

Let $\mathbf{A} \leq \mathbf{B}$ and \mathcal{K} a class. Say that $b \in \mathbf{B}$ is **dominated** by \mathbf{A} w.r.t. \mathcal{K} if for all $\mathbf{C} \in \mathcal{K}$, and all $h, h' : \mathbf{B} \rightarrow \mathbf{C}$

$$h|_A = h'|_A \text{ implies } h(b) = h'(b).$$

Dominions

Let $\mathbf{A} \leq \mathbf{B}$ and \mathcal{K} a class. Say that $b \in \mathbf{B}$ is **dominated** by \mathbf{A} w.r.t. \mathcal{K} if for all $\mathbf{C} \in \mathcal{K}$, and all $h, h' : \mathbf{B} \rightarrow \mathbf{C}$

$$h|_A = h'|_A \text{ implies } h(b) = h'(b).$$

The **dominion** of \mathbf{A} in \mathbf{B} w.r.t. \mathcal{K} , $\text{dom}_{\mathcal{K}}^{\mathbf{B}} \mathbf{A}$, is the set of all such b .

Dominions


Let $\mathbf{A} \leq \mathbf{B}$ and \mathcal{K} a class. Say that $b \in \mathbf{B}$ is **dominated** by \mathbf{A} w.r.t. \mathcal{K} if for all $\mathbf{C} \in \mathcal{K}$, and all $h, h' : \mathbf{B} \rightarrow \mathbf{C}$

$$h|_A = h'|_A \text{ implies } h(b) = h'(b).$$

The **dominion** of \mathbf{A} in \mathbf{B} w.r.t. \mathcal{K} , $\text{dom}_{\mathcal{K}}^{\mathbf{B}} \mathbf{A}$, is the set of all such b .

Example

Let \mathcal{D}_{01} be the variety of bounded distributive lattices.



Dominions


Let $\mathbf{A} \leq \mathbf{B}$ and \mathcal{K} a class. Say that $b \in \mathbf{B}$ is **dominated** by \mathbf{A} w.r.t. \mathcal{K} if for all $\mathbf{C} \in \mathcal{K}$, and all $h, h' : \mathbf{B} \rightarrow \mathbf{C}$

$$h|_A = h'|_A \text{ implies } h(b) = h'(b).$$

The **dominion** of \mathbf{A} in \mathbf{B} w.r.t. \mathcal{K} , $\text{dom}_{\mathcal{K}}^{\mathbf{B}} \mathbf{A}$, is the set of all such b .

Example

Let \mathcal{D}_{01} be the variety of bounded distributive lattices.



Observe that $b \in \text{dom}_{\mathcal{D}_{01}}^{\mathbf{B}} \mathbf{A}$.

Dominions in \mathcal{D}_{01}

Let \mathcal{B} be the class of boolean algebras. Given $\mathbf{A} \in \mathcal{D}_{01}$, there is a unique $\bar{\mathbf{A}} \in \mathcal{B}$ such that:

- ▶ $\mathbf{A} \leq \bar{\mathbf{A}}|_{\mathcal{D}_{01}}$
- ▶ \mathbf{A} generates $\bar{\mathbf{A}}$.

Dominions in \mathcal{D}_{01}

Let \mathcal{B} be the class of boolean algebras. Given $\mathbf{A} \in \mathcal{D}_{01}$, there is a unique $\bar{\mathbf{A}} \in \mathcal{B}$ such that:

- ▶ $\mathbf{A} \leq \bar{\mathbf{A}}|_{\mathcal{D}_{01}}$
- ▶ A generates $\bar{\mathbf{A}}$.

Theorem (Wasserman 2001)

Let $\mathbf{A} \leq \mathbf{B} \in \mathcal{D}_{01}$, and let $b \in B$. T.f.a.e.:

- ▶ $b \in \text{dom}_{\mathcal{D}_{01}}^{\mathbf{B}} \mathbf{A}$
- ▶ b is in the subalgebra of $\bar{\mathbf{B}}$ generated by A .

Discriminator varieties

The (quaternary) **discriminator** on a set A is the function $d_A : A^4 \rightarrow A$ defined by

$$d_A(x, y, z, w) = \begin{cases} z & \text{if } x = y \\ w & \text{if } x \neq y \end{cases}$$

Discriminator varieties

The (quaternary) **discriminator** on a set A is the function $d_A : A^4 \rightarrow A$ defined by

$$d_A(x, y, z, w) = \begin{cases} z & \text{if } x = y \\ w & \text{if } x \neq y \end{cases}$$

A variety \mathcal{V} is a **discriminator variety** if there are a class \mathcal{M} and a term $t(x, y, z, w)$ such that

Discriminator varieties

The (quaternary) **discriminator** on a set A is the function $d_A : A^4 \rightarrow A$ defined by

$$d_A(x, y, z, w) = \begin{cases} z & \text{if } x = y \\ w & \text{if } x \neq y \end{cases}$$

A variety \mathcal{V} is a **discriminator variety** if there are a class \mathcal{M} and a term $t(x, y, z, w)$ such that

$$\blacktriangleright \mathcal{V} = \text{HSP}(\mathcal{M})$$

Discriminator varieties

The (quaternary) **discriminator** on a set A is the function $d_A : A^4 \rightarrow A$ defined by

$$d_A(x, y, z, w) = \begin{cases} z & \text{if } x = y \\ w & \text{if } x \neq y \end{cases}$$

A variety \mathcal{V} is a **discriminator variety** if there are a class \mathcal{M} and a term $t(x, y, z, w)$ such that

- ▶ $\mathcal{V} = \text{HSP}(\mathcal{M})$
- ▶ $t^{\mathbf{A}} = d_A$ for each $\mathbf{A} \in \mathcal{M}$.

Discriminator varieties

The (quaternary) **discriminator** on a set A is the function $d_A : A^4 \rightarrow A$ defined by

$$d_A(x, y, z, w) = \begin{cases} z & \text{if } x = y \\ w & \text{if } x \neq y \end{cases}$$

A variety \mathcal{V} is a **discriminator variety** if there are a class \mathcal{M} and a term $t(x, y, z, w)$ such that

- ▶ $\mathcal{V} = \text{HSP}(\mathcal{M})$
- ▶ $t^{\mathbf{A}} = d_A$ for each $\mathbf{A} \in \mathcal{M}$.

Examples:

- ▶ Boolean algebras,

Discriminator varieties

The (quaternary) **discriminator** on a set A is the function $d_A : A^4 \rightarrow A$ defined by

$$d_A(x, y, z, w) = \begin{cases} z & \text{if } x = y \\ w & \text{if } x \neq y \end{cases}$$

A variety \mathcal{V} is a **discriminator variety** if there are a class \mathcal{M} and a term $t(x, y, z, w)$ such that

- ▶ $\mathcal{V} = \text{HSP}(\mathcal{M})$
- ▶ $t^{\mathbf{A}} = d_A$ for each $\mathbf{A} \in \mathcal{M}$.

Examples:

- ▶ Boolean algebras,
- ▶ Varieties of rings generated by finitely many finite fields,

Discriminator varieties

The (quaternary) **discriminator** on a set A is the function $d_A : A^4 \rightarrow A$ defined by

$$d_A(x, y, z, w) = \begin{cases} z & \text{if } x = y \\ w & \text{if } x \neq y \end{cases}$$

A variety \mathcal{V} is a **discriminator variety** if there are a class \mathcal{M} and a term $t(x, y, z, w)$ such that

- ▶ $\mathcal{V} = \mathbf{HSP}(\mathcal{M})$
- ▶ $t^{\mathbf{A}} = d_A$ for each $\mathbf{A} \in \mathcal{M}$.

Examples:

- ▶ Boolean algebras,
- ▶ Varieties of rings generated by finitely many finite fields,
- ▶ Monadic algebras.

Filtral quasivarieties

A quasivariety \mathcal{Q} is **filtral** if

- ▶ \mathcal{Q} is congruence distributive
- ▶ \mathcal{Q} is semisimple
- ▶ \mathcal{Q}_S is a universal class.

Filtral quasivarieties

A quasivariety \mathcal{Q} is **filtral** if

- ▶ \mathcal{Q} is congruence distributive
- ▶ \mathcal{Q} is semisimple
- ▶ \mathcal{Q}_S is a universal class.

Examples:

- ▶ Discriminator varieties,

Filtral quasivarieties

A quasivariety \mathcal{Q} is **filtral** if

- ▶ \mathcal{Q} is congruence distributive
- ▶ \mathcal{Q} is semisimple
- ▶ \mathcal{Q}_S is a universal class.

Examples:

- ▶ Discriminator varieties,
- ▶ Distributive lattices,

Filtral quasivarieties

A quasivariety \mathcal{Q} is **filtral** if

- ▶ \mathcal{Q} is congruence distributive
- ▶ \mathcal{Q} is semisimple
- ▶ \mathcal{Q}_S is a universal class.

Examples:

- ▶ Discriminator varieties,
- ▶ Distributive lattices,
- ▶ De Morgan algebras.

Filtral quasivarieties

A quasivariety \mathcal{Q} is **filtral** if

- ▶ \mathcal{Q} is congruence distributive
- ▶ \mathcal{Q} is semisimple
- ▶ \mathcal{Q}_S is a universal class.

Examples:

- ▶ Discriminator varieties,
- ▶ Distributive lattices,
- ▶ De Morgan algebras.

Theorem (C. & Vaggione 2012)

A quasivariety \mathcal{Q} is filtral iff there are a class \mathcal{M} and a conjunction of atomic formulas $\alpha(x, y, z, w, u)$ such that

Filtral quasivarieties

A quasivariety \mathcal{Q} is **filtral** if

- ▶ \mathcal{Q} is congruence distributive
- ▶ \mathcal{Q} is semisimple
- ▶ \mathcal{Q}_S is a universal class.

Examples:

- ▶ Discriminator varieties,
- ▶ Distributive lattices,
- ▶ De Morgan algebras.

Theorem (C. & Vaggione 2012)

A quasivariety \mathcal{Q} is filtral iff there are a class \mathcal{M} and a conjunction of atomic formulas $\alpha(x, y, z, w, u)$ such that

- ▶ $\mathcal{Q} = \text{ISPP}_U(\mathcal{M})$,
- ▶ For each $\mathbf{A} \in \mathcal{M}$ we have
 $\mathbf{A} \models \alpha(x, y, z, w, u) \leftrightarrow d_{\mathbf{A}}(x, y, z, w) = u$.

Existentially closed structures and the AP

Let \mathcal{M} be a class of structures.

Existentially closed structures and the AP

Let \mathcal{M} be a class of structures.

- ▶ \mathbf{A} is **existentially closed** (ec) in \mathcal{M} if for all $\mathbf{B} \in \mathcal{M}$ with $\mathbf{A} \leq \mathbf{B}$, every existential formula $\varphi(\vec{x})$, and all $\vec{a} \in A$ we have

$$\mathbf{B} \models \varphi(\vec{a}) \implies \mathbf{A} \models \varphi(\vec{a}).$$

Existentially closed structures and the AP

Let \mathcal{M} be a class of structures.

- ▶ \mathbf{A} is **existentially closed** (ec) in \mathcal{M} if for all $\mathbf{B} \in \mathcal{M}$ with $\mathbf{A} \leq \mathbf{B}$, every existential formula $\varphi(\bar{x})$, and all $\bar{a} \in A$ we have

$$\mathbf{B} \models \varphi(\bar{a}) \implies \mathbf{A} \models \varphi(\bar{a}).$$

Let $\mathcal{M}_{\text{ec}} := \{\mathbf{A} : \mathbf{A} \text{ is ec in } \mathcal{M}\}.$

Existentially closed structures and the AP

Let \mathcal{M} be a class of structures.

- ▶ \mathbf{A} is **existentially closed** (ec) in \mathcal{M} if for all $\mathbf{B} \in \mathcal{M}$ with $\mathbf{A} \leq \mathbf{B}$, every existential formula $\varphi(\bar{x})$, and all $\bar{a} \in A$ we have

$$\mathbf{B} \models \varphi(\bar{a}) \implies \mathbf{A} \models \varphi(\bar{a}).$$

Let $\mathcal{M}_{\text{ec}} := \{\mathbf{A} : \mathbf{A} \text{ is ec in } \mathcal{M}\}.$

- ▶ \mathcal{M} has the **amalgamation property** (AP) if

Existentially closed structures and the AP

Let \mathcal{M} be a class of structures.

- ▶ \mathbf{A} is **existentially closed** (ec) in \mathcal{M} if for all $\mathbf{B} \in \mathcal{M}$ with $\mathbf{A} \leq \mathbf{B}$, every existential formula $\varphi(\bar{x})$, and all $\bar{a} \in A$ we have

$$\mathbf{B} \models \varphi(\bar{a}) \implies \mathbf{A} \models \varphi(\bar{a}).$$

Let $\mathcal{M}_{\text{ec}} := \{\mathbf{A} : \mathbf{A} \text{ is ec in } \mathcal{M}\}.$

- ▶ \mathcal{M} has the **amalgamation property** (AP) if (see blackboard \downarrow)

Dominions are generated by definable functions

Let \mathcal{Q} be a filtral quasivariety and \mathcal{M} its class of simple members.

Dominions are generated by definable functions

Let \mathcal{Q} be a filtral quasivariety and \mathcal{M} its class of simple members.

Theorem

Suppose \mathcal{M} has the AP and \mathcal{M}_{ec} is axiomatizable. Let $\mathbf{A} \leq \mathbf{B} \in \mathcal{Q}$ and $b \in B$, the following are equivalent:

- ▶ $b \in \text{dom}_{\mathbf{B}}^{\mathcal{Q}} \mathbf{A}$

Dominions are generated by definable functions

Let \mathcal{Q} be a filtral quasivariety and \mathcal{M} its class of simple members.

Theorem

Suppose \mathcal{M} has the AP and \mathcal{M}_{ec} is axiomatizable. Let $\mathbf{A} \leq \mathbf{B} \in \mathcal{Q}$ and $b \in B$, the following are equivalent:

- ▶ $b \in \text{dom}_{\mathbf{B}}^{\mathcal{Q}} \mathbf{A}$
- ▶ There are a conjunction of equations $\delta(\bar{x}, y)$ and $\bar{a} \in A$ such that:
 - ▶ $\mathbf{B} \models \delta(\bar{a}, b)$

Dominions are generated by definable functions

Let \mathcal{Q} be a filtral quasivariety and \mathcal{M} its class of simple members.

Theorem

Suppose \mathcal{M} has the AP and \mathcal{M}_{ec} is axiomatizable. Let $\mathbf{A} \leq \mathbf{B} \in \mathcal{Q}$ and $b \in B$, the following are equivalent:

- ▶ $b \in \text{dom}_{\mathbf{B}}^{\mathcal{Q}} \mathbf{A}$
- ▶ There are a conjunction of equations $\delta(\bar{x}, y)$ and $\bar{a} \in A$ such that:
 - ▶ $\mathbf{B} \models \delta(\bar{a}, b)$
 - ▶ $\mathcal{M}_{\text{ec}} \models \forall \bar{x} \exists ! y \delta(\bar{x}, y)$.

The saturated expansion

Let

$$\Delta := \{\delta(\bar{x}, y) : \mathcal{M}_{ec} \models \forall \bar{x} \exists! y \delta(\bar{x}, y)\}.$$

The saturated expansion

Let

$$\Delta := \{\delta(\bar{x}, y) : \mathcal{M}_{ec} \models \forall \bar{x} \exists! y \delta(\bar{x}, y)\}.$$

For each $\delta(x_1, \dots, x_n, y) \in \Delta$ let f_δ be a new n -ary function symbol and let

$$\mathcal{L}_\Delta := \mathcal{L}_\psi \cup \{f_\delta : \delta \in \Delta\},$$

The saturated expansion

Let

$$\Delta := \{\delta(\bar{x}, y) : \mathcal{M}_{ec} \models \forall \bar{x} \exists! y \delta(\bar{x}, y)\}.$$

For each $\delta(x_1, \dots, x_n, y) \in \Delta$ let f_δ be a new n -ary function symbol and let

$$\mathcal{L}_\Delta := \mathcal{L}_\psi \cup \{f_\delta : \delta \in \Delta\},$$

Define

$$\blacktriangleright \Gamma := \{\forall \bar{x} \delta(\bar{x}, f_\delta(\bar{x})) : \delta \in \Delta\} \cup \{\text{axioms for } \mathcal{Q}\},$$

The saturated expansion

Let

$$\Delta := \{\delta(\bar{x}, y) : \mathcal{M}_{ec} \models \forall \bar{x} \exists! y \delta(\bar{x}, y)\}.$$

For each $\delta(x_1, \dots, x_n, y) \in \Delta$ let f_δ be a new n -ary function symbol and let

$$\mathcal{L}_\Delta := \mathcal{L}_\psi \cup \{f_\delta : \delta \in \Delta\},$$

Define

- ▶ $\Gamma := \{\forall \bar{x} \delta(\bar{x}, f_\delta(\bar{x})) : \delta \in \Delta\} \cup \{\text{axioms for } \mathcal{Q}\},$
- ▶ $\mathcal{Q}^\Delta := \text{Mod}(\Gamma).$

Characterization of dominions in filtral quasivarieties

Theorem

- ▶ \mathcal{Q}^Δ is a discriminator variety.

Characterization of dominions in filtral quasivarieties

Theorem

- ▶ \mathcal{Q}^Δ is a discriminator variety.
- ▶ Dominions are trivial in \mathcal{Q}^Δ .

Characterization of dominions in filtral quasivarieties

Theorem

- ▶ \mathcal{Q}^Δ is a discriminator variety.
- ▶ Dominions are trivial in \mathcal{Q}^Δ .
- ▶ For every $\mathbf{A} \in \mathcal{Q}$ there is a unique $\bar{\mathbf{A}} \in \mathcal{Q}^\Delta$ such that $\mathbf{A} \leq \bar{\mathbf{A}}|_{\mathcal{Q}}$ and \mathbf{A} generates $\bar{\mathbf{A}}$.

Characterization of dominions in filtral quasivarieties

Theorem

- ▶ \mathcal{Q}^Δ is a discriminator variety.
- ▶ Dominions are trivial in \mathcal{Q}^Δ .
- ▶ For every $\mathbf{A} \in \mathcal{Q}$ there is a unique $\bar{\mathbf{A}} \in \mathcal{Q}^\Delta$ such that $\mathbf{A} \leq \bar{\mathbf{A}}|_{\mathcal{Q}}$ and A generates $\bar{\mathbf{A}}$.
- ▶ $\text{dom}_{\mathbf{B}}^{\mathcal{Q}} \mathbf{A} = \langle A \rangle^{\bar{\mathbf{B}}} \cap B$ for all $\mathbf{A} \leq \mathbf{B} \in \mathcal{Q}$.

Thank you!

