

On the Logical Implications of Proof Forms

Raheleh Jalali

(joint work with Amir Akbar Tabatabai)

Institute of Mathematics, Czech Academy of Sciences

Logic Colloquium, Prague

15 August, 2019

Motivation

Proof systems play a crucial role in proof theory, from consistency proofs and proof mining techniques to the characterization of admissible rules.

Motivation

Proof systems play a crucial role in proof theory, from consistency proofs and proof mining techniques to the characterization of admissible rules. Our goal: study proof systems in a generic manner.

Motivation

Proof systems play a crucial role in proof theory, from consistency proofs and proof mining techniques to the characterization of admissible rules. Our goal: study proof systems in a generic manner.

Question

Is it possible to prove that some logics do not have a “nice” proof system?

Motivation

Proof systems play a crucial role in proof theory, from consistency proofs and proof mining techniques to the characterization of admissible rules. Our goal: study proof systems in a generic manner.

Question

Is it possible to prove that some logics do not have a “nice” proof system?

This problem has three sides:

- Formalizing nice proof systems;
- considering their corresponding logics;
- finding an invariant, i.e., a property that the logic of a nice proof system enjoys.

Prove almost all logics in a certain given class do not enjoy that property.

Theorem (Iemhoff [5])

If a *super-intuitionistic* logic has a terminating proof system consisting of *focused rules* and *focused axioms*, it has the uniform interpolation property.

Theorem (Iemhoff [5])

If a *super-intuitionistic* logic has a terminating proof system consisting of *focused rules* and *focused axioms*, it has the uniform interpolation property.

- ▶ *Focused axioms* are a modest generalization of the axioms of **LJ**.

Theorem (Iemhoff [5])

If a *super-intuitionistic* logic has a terminating proof system consisting of *focused rules* and *focused axioms*, it has the uniform interpolation property.

- ▶ *Focused axioms* are a modest generalization of the axioms of **LJ**.
- ▶ A *focused rule* is a rule with one main formula in its consequence such that the rule respects both the side of this main formula and the occurrence of atoms in it.

Theorem (Iemhoff [5])

If a *super-intuitionistic* logic has a terminating proof system consisting of *focused rules* and *focused axioms*, it has the uniform interpolation property.

- ▶ *Focused axioms* are a modest generalization of the axioms of **LJ**.
- ▶ A *focused rule* is a rule with one main formula in its consequence such that the rule respects both the side of this main formula and the occurrence of atoms in it. Example: Conjunction and disjunction rules are focused; implication rules are not.

Theorem (Iemhoff [5])

If a *super-intuitionistic* logic has a terminating proof system consisting of *focused rules* and *focused axioms*, it has the uniform interpolation property.

- ▶ *Focused axioms* are a modest generalization of the axioms of **LJ**.
- ▶ A *focused rule* is a rule with one main formula in its consequence such that the rule respects both the side of this main formula and the occurrence of atoms in it. Example: Conjunction and disjunction rules are focused; implication rules are not.

- Nice proof systems are focused proof systems;
- corresponding logics are super-intuitionistic;
- the invariant is uniform interpolation.

Only seven super-intuitionistic logics have uniform interpolation.

Our Contribution

- We will present a second approximation for *nice* proof systems.
- Our candidate for natural well-behaved sequent-style rules is *semi-analytic* rules (focused rules with no side preserving condition).
- It covers a vast variety of rules: focused rules, implication rules, non-context sharing rules in substructural logics and so many others. We also consider the usual modal rules K and D .

Then, we show:

Main Result (informal.)

Theorem (Akbar Tabatabai, J.)

- (i) If a *sufficiently strong sub-structural logic* has a sequent-style proof system only consisting of *semi-analytic* rules and focused axioms, it has the Craig interpolation property. As a result, many substructural logics and all super-intuitionistic logics, except seven of them, do not have a sequent calculus of the mentioned form.
- (ii) If a *sufficiently strong sub-structural logic* has a **terminating** sequent-style proof system only consisting of semi-analytic rules and focused axioms, it has the uniform interpolation property. Consequently, **K4** and **S4** do not have a terminating sequent calculus of the mentioned form.

The theorem provides a uniform, proof theoretical and modular method to prove Craig and uniform interpolation.

Craig interpolation

We say a logic L has Craig interpolation property if for any formulas ϕ and ψ if $L \vdash \phi \rightarrow \psi$, then there exists formula θ such that $L \vdash \phi \rightarrow \theta$ and $L \vdash \theta \rightarrow \psi$ and $V(\theta) \subseteq V(\phi) \cap V(\psi)$.

Craig interpolation

We say a logic L has Craig interpolation property if for any formulas ϕ and ψ if $L \vdash \phi \rightarrow \psi$, then there exists formula θ such that $L \vdash \phi \rightarrow \theta$ and $L \vdash \theta \rightarrow \psi$ and $V(\theta) \subseteq V(\phi) \cap V(\psi)$.

Uniform interpolation

We say a logic L has the uniform interpolation property if for any formula ϕ and any atomic formula p , there are two p -free formulas, the p -pre-interpolant, $\forall p\phi$ and the p -post-interpolant $\exists p\phi$, such that $V(\exists p\phi) \subseteq V(\phi)$ and $V(\forall p\phi) \subseteq V(\phi)$ and

- (i) $L \vdash \forall p\phi \rightarrow \phi$,
- (ii) For any p -free formula ψ if $L \vdash \psi \rightarrow \phi$ then $L \vdash \psi \rightarrow \forall p\phi$,
- (iii) $L \vdash \phi \rightarrow \exists p\phi$, and
- (iv) For any p -free formula ψ if $L \vdash \phi \rightarrow \psi$ then $L \vdash \exists p\phi \rightarrow \psi$.

Terminating calculus: there is an order on the sequents...

Basic Sub-structural Logics

$$\overline{\phi \Rightarrow \phi} \quad \overline{\Rightarrow 1} \quad \overline{0 \Rightarrow} \quad \overline{\Gamma \Rightarrow \top, \Delta} \quad \overline{\Gamma, \perp \Rightarrow \Delta}$$

$$\frac{\Gamma \Rightarrow \Delta}{\Gamma, 1 \Rightarrow \Delta} L1 \quad \frac{\Gamma \Rightarrow \Delta}{\Gamma \Rightarrow 0, \Delta} R0$$

$$\frac{\Gamma, \phi \Rightarrow \Delta}{\Gamma, \phi \wedge \psi \Rightarrow \Delta} L\wedge \quad \frac{\Gamma \Rightarrow \phi, \Delta \quad \Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \wedge \psi, \Delta} R\wedge$$

$$\frac{\Gamma, \phi \Rightarrow \Delta \quad \Gamma, \psi \Rightarrow \Delta}{\Gamma, \phi \vee \psi \Rightarrow \Delta} L\vee \quad \frac{\Gamma \Rightarrow \phi, \Delta}{\Gamma \Rightarrow \phi \vee \psi, \Delta} R\vee \quad \frac{\Gamma \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \vee \psi, \Delta} R\vee$$

$$\frac{\Gamma, \phi, \psi \Rightarrow \Delta}{\Gamma, \phi * \psi \Rightarrow \Delta} L* \quad \frac{\Gamma \Rightarrow \phi, \Delta \quad \Sigma \Rightarrow \psi, \Lambda}{\Gamma, \Sigma \Rightarrow \phi * \psi, \Delta, \Lambda} R*$$

$$\frac{\Gamma \Rightarrow \phi, \Delta \quad \Sigma, \psi \Rightarrow \Lambda}{\Gamma, \Sigma, \phi \rightarrow \psi \Rightarrow \Delta, \Lambda} L\rightarrow \quad \frac{\Gamma, \phi \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \rightarrow \psi, \Delta} R\rightarrow$$

Basic Sub-structural Logics

- The system consisting of the single-conclusion version of all of the above-mentioned rules is \mathbf{FL}_e .

Basic Sub-structural Logics

- The system consisting of the single-conclusion version of all of the above-mentioned rules is \mathbf{FL}_e .
- In the multi-conclusion case define \mathbf{CFL}_e with the same rules as \mathbf{FL}_e , this time in their full multi-conclusion version and add $+$ to the language and the following rules to the system:

$$\frac{\Gamma, \phi \Rightarrow \Delta \quad \Sigma, \psi \Rightarrow \Lambda}{\Gamma, \Sigma, \phi + \psi \Rightarrow \Delta, \Lambda} L+ \quad \frac{\Gamma \Rightarrow \phi, \psi, \Delta}{\Gamma \Rightarrow \phi + \psi, \Delta} R+$$

Structural Rules

Weakening rules:

$$\frac{\Gamma \Rightarrow \Delta}{\Gamma, \phi \Rightarrow \Delta} Lw \quad \frac{\Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \phi, \Delta} Rw$$

Contraction rules:

$$\frac{\Gamma, \phi, \phi \Rightarrow \Delta}{\Gamma, \phi \Rightarrow \Delta} Lc \quad \frac{\Gamma \Rightarrow \phi, \phi, \Delta}{\Gamma \Rightarrow \phi, \Delta} Rc$$

Weakening rules:

$$\frac{\Gamma \Rightarrow \Delta}{\Gamma, \phi \Rightarrow \Delta} Lw \quad \frac{\Gamma \Rightarrow \Delta}{\Gamma \Rightarrow \phi, \Delta} Rw$$

Contraction rules:

$$\frac{\Gamma, \phi, \phi \Rightarrow \Delta}{\Gamma, \phi \Rightarrow \Delta} Lc \quad \frac{\Gamma \Rightarrow \phi, \phi, \Delta}{\Gamma \Rightarrow \phi, \Delta} Rc$$

- $\mathbf{FL}_{ew} = \mathbf{FL}_e + (Lw) + (Rw)$,
- $\mathbf{FL}_{ec} = \mathbf{FL}_e + (Lc)$,
- $\mathbf{CFL}_{ew} = \mathbf{CFL}_e + (Lw) + (Rw)$,
- $\mathbf{CFL}_{ec} = \mathbf{CFL}_e + (Lc) + (Rc)$.

Semi-analytic rules: Single-conclusion

- *Left semi-analytic rule:*

$$\frac{\langle\langle\Pi_j, \bar{\psi}_{js} \Rightarrow \bar{\theta}_{js}\rangle_s\rangle_j \quad \langle\langle\Gamma_i, \bar{\phi}_{ir} \Rightarrow \Delta_i\rangle_r\rangle_i}{\Pi_1, \dots, \Pi_m, \Gamma_1, \dots, \Gamma_n, \phi \Rightarrow \Delta_1, \dots, \Delta_n}$$

where Π_j , Γ_i and Δ_i 's are meta-multiset variables and

$$\bigcup_{i,r} V(\bar{\phi}_{ir}) \cup \bigcup_{j,s} V(\bar{\psi}_{js}) \cup \bigcup_{j,s} V(\bar{\theta}_{js}) \subseteq V(\phi)$$

Semi-analytic rules: Single-conclusion

- *Left semi-analytic rule:*

$$\frac{\langle\langle\Pi_j, \bar{\psi}_{js} \Rightarrow \bar{\theta}_{js}\rangle_s\rangle_j \quad \langle\langle\Gamma_i, \bar{\phi}_{ir} \Rightarrow \Delta_i\rangle_r\rangle_i}{\Pi_1, \dots, \Pi_m, \Gamma_1, \dots, \Gamma_n, \phi \Rightarrow \Delta_1, \dots, \Delta_n}$$

where Π_j , Γ_i and Δ_i 's are meta-multiset variables and

$$\bigcup_{i,r} V(\bar{\phi}_{ir}) \cup \bigcup_{j,s} V(\bar{\psi}_{js}) \cup \bigcup_{j,s} V(\bar{\theta}_{js}) \subseteq V(\phi)$$

- *Right semi-analytic rule:*

$$\frac{\langle\langle\Gamma_i, \bar{\phi}_{ir} \Rightarrow \bar{\psi}_{ir}\rangle_r\rangle_i}{\Gamma_1, \dots, \Gamma_n \Rightarrow \phi}$$

Semi-analytic rules: Multi-conclusion

- *Left multi-conclusion semi-analytic rule:*

$$\frac{\langle\langle\Gamma_i, \bar{\phi}_{ir} \Rightarrow \bar{\psi}_{ir}, \Delta_i\rangle_r\rangle_i}{\Gamma_1, \dots, \Gamma_n, \phi \Rightarrow \Delta_1, \dots, \Delta_n}$$

- *Right multi-conclusion semi-analytic rule:*

$$\frac{\langle\langle\Gamma_i, \bar{\phi}_{ir} \Rightarrow \bar{\psi}_{ir}, \Delta_i\rangle_r\rangle_i}{\Gamma_1, \dots, \Gamma_n \Rightarrow \phi, \Delta_1, \dots, \Delta_n}$$

Semi-analytic modal rules

A rule is called *modal semi-analytic* if it has one of the following forms:

$$\frac{\Gamma \Rightarrow \phi}{\Box \Gamma \Rightarrow \Box \phi} K \quad \frac{\Gamma \Rightarrow}{\Box \Gamma \Rightarrow} D$$

with the conditions that first, Γ is a meta-multiset variable and secondly whenever the rule (D) is present, the rule (K) must be present, as well.

Example

A generic example of a left semi-analytic rule is the following:

$$\frac{\Gamma, \phi_1, \phi_2 \Rightarrow \psi \quad \Gamma, \theta \Rightarrow \eta \quad \Pi, \mu_1, \mu_2, \mu_3 \Rightarrow \Delta}{\Gamma, \Pi, \alpha \Rightarrow \Delta}$$

where

$$V(\phi_1, \phi_2, \psi, \theta, \eta, \mu_1, \mu_2, \mu_3) \subseteq V(\alpha)$$

Example

The following rules are semi-analytic:

- ▶ the usual conjunction, disjunction and implication rules for **IPC**;
- ▶ all the rules in sub-structural logic **FL_e**, weakening and contraction rules;
- ▶ the following rules for exponentials in linear logic:

$$\frac{\Gamma, !\phi, !\phi \Rightarrow \Delta}{\Gamma, !\phi \Rightarrow \Delta} \quad \frac{\Gamma \Rightarrow \Delta}{\Gamma, !\phi \Rightarrow \Delta}$$

Example

- ▶ The cut rule; since it does not meet the variable occurrence condition.
- ▶ the following rule in the calculus of **KC**:

$$\frac{\Gamma, \phi \Rightarrow \psi, \Delta}{\Gamma \Rightarrow \phi \rightarrow \psi, \Delta}$$

in which Δ should consist of negation formulas is not a multi-conclusion semi-analytic rule, simply because the context is not free for all possible substitutions.

Focused axioms

A sequent is called a *focused axiom* if it has the following form:

- (1) $(\phi \Rightarrow \phi)$
- (2) $(\Rightarrow \bar{\alpha})$
- (3) $(\bar{\beta} \Rightarrow)$
- (4) $(\Gamma, \bar{\phi} \Rightarrow \Delta)$
- (5) $(\Gamma \Rightarrow \bar{\phi}, \Delta)$

where Γ and Δ are meta-multiset variables and in (2) – (5) the variables in any pair of elements in $\bar{\alpha}$ or $\bar{\beta}$ or $\bar{\phi}$ are equal.

Example

It is easy to see that the axioms given in the preliminaries are examples of focused axioms. Here are some more examples:

$$\neg 1 \Rightarrow \quad , \quad \Rightarrow \neg 0$$

$$\phi, \neg\phi \Rightarrow \quad , \quad \Rightarrow \phi, \neg\phi$$

$$\Gamma, \neg\top \Rightarrow \Delta \quad , \quad \Gamma \Rightarrow \Delta, \neg\perp$$

Main Result (formal.)

Theorem

- (i) *If $\mathbf{FL}_e \subseteq L$, and L has a (terminating) single-conclusion sequent calculus consisting of semi-analytic rules and focused axioms, then L has Craig (uniform) interpolation.*
- (ii) *If $\mathbf{IPC} \subseteq L$ and L has a single-conclusion sequent calculus consisting of semi-analytic rules and focused axioms, then L has Craig interpolation.*
- (iii) *If $\mathbf{CFL}_e \subseteq L$, and L has a (terminating) multi-conclusion sequent calculus consisting of semi-analytic rules and focused axioms, then L has Craig (uniform) interpolation.*

As a positive application we have the following:

Corollary

The logics \mathbf{FL}_e , \mathbf{FL}_{ew} , \mathbf{CFL}_e , \mathbf{CFL}_{ew} , \mathbf{CPC} , and their \mathbf{K} and \mathbf{KD} modal versions have the uniform interpolation property.

Positive Application

As a positive application we have the following:

Corollary

The logics \mathbf{FL}_e , \mathbf{FL}_{ew} , \mathbf{CFL}_e , \mathbf{CFL}_{ew} , \mathbf{CPC} , and their \mathbf{K} and \mathbf{KD} modal versions have the uniform interpolation property.

Proof.

The usual sequent calculi for these logics consist of some suitable variants of semi-analytic rules and modal rules. □

Corollary

*None of the following logics can have a **nice** proof system:*

- ▶ *Many substructural logics (\mathbf{L}_n , \mathbf{L}_∞ , \mathbf{R} , \mathbf{BL} , \dots);*
- ▶ *Almost all super-intuitionistic logics (except at most seven of them);*
- ▶ *Almost all extensions of **S4** (except at most thirty seven of them);*

Thank you!

-  Bílková, M. Uniform interpolation and propositional quantifiers in modal logics. *Studia Logica*, 85(1):1-31, 2007.
-  Marchioni E. and Metcalfe G. Craig interpolation for semilinear substructural logics, *Mathematical Logic Quarterly*, Volume 58, Issue 6 November 2012 Pages 468-481.
-  Ghilardi, S. and Zawadowski, M. Sheaves, games, and model completions: a categorical approach to non-classical propositional logics, volume 14. Springer Science and Business Media, 2013.
-  Alizadeh, M., Derakhshan, F. and Ono, H. Uniform interpolation in substructural logics. *The Review of Symbolic Logic*, 2014.
-  Iemhoff, R. Uniform interpolation and the existence of sequent calculi, 2017.
-  Akbar Tabatabai A., Jalali R. *Universal Proof Theory: Semi-analytic Rules and Craig Interpolation*, 2019.